

polyCub: An R package for Integration over Polygons

Sebastian Meyer¹

¹ Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), Erlangen, Germany

DOI: [10.21105/joss.01056](https://doi.org/10.21105/joss.01056)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Submitted: 18 October 2018

Published: 12 February 2019

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC-BY](#)).

Summary

The integral of a continuously differentiable function $f(x, y)$ over a domain $W \subset \mathbb{R}^2$ can be approximated using an n -point cubature rule of the form

$$\iint_W f(x, y) \, dx \, dy \approx \sum_{i=1}^n w_i f(x_i, y_i),$$

i.e., a weighted sum of function values at an appropriate set of nodes. In the special but common case of integration along the axes, i.e., $W = (x_l, x_u) \times (y_l, y_u)$, the domain is *rectangular*. Several software packages implement numerical integration over such rectangles (or *hypercubes* in higher dimensions), for example, the `Cuba` (Hahn, 2005) or `cubature` (Johnson, 2017) libraries, which are both interfaced from the `cubature` package (Narasimhan, Johnson, Hahn, Bouvier, & Kiêu, 2018) in R (R Core Team, 2018).

In spatial statistics, however, the domains of interest typically correspond to geographic regions (administrative districts, lakes, etc.), which are described by *polygons*. Solving integrals over such complex domains requires specialized cubature methods, thus the R package `polyCub`. A simple graphical summary of the purpose of `polyCub` is given by its logo (see below).

`polyCub` implements the following methods for numerical integration over polygons:

- General-purpose *product Gauss cubature* (Sommariva & Vianello, 2007)
- Simple *two-dimensional midpoint rule* via `spatstat` (Baddeley & Turner, 2005)
- Adaptive cubature for *radially symmetric functions* $f(x, y) = f_r(\|(x - x_0, y - y_0)\|)$ via integration along the polygon boundary (Meyer & Held, 2014, Supplement B)
- Accurate (but slow) integration of the *bivariate Gaussian density* based on polygon triangulation (Abramowitz & Stegun, 1972, Section 26.9, Example 9)



Figure 1: polyCub: cubature over polygonal domains.

Usage

The R package `polyCub` is released on the Comprehensive R Archive Network ([CRAN](#)) and can thus be easily installed using `install.packages("polyCub")` in R. After that, the basic usage is

```
library("polyCub")
polyCub(polyregion, f)
```

where `polyregion` is the integration domain and `f` is the integrand. Details are given in

```
vignette("polyCub")
```

which exemplifies the implemented cubature methods by solving the integral displayed in the package logo.

`polyCub` is currently used by at least two other R packages: in `surveillance`, to evaluate the likelihood of self-exciting spatio-temporal point process models for infectious disease spread (Meyer, Held, & Höhle, 2017), and in `rase`, to integrate bivariate Gaussian densities for phylogeographic analyses (Quintero, Keil, Jetz, & Crawford, 2015).

References

- Abramowitz, M., & Stegun, I. A. (Eds.). (1972). *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. National bureau of standards applied mathematics series. New York: Dover Publications.
- Baddeley, A., & Turner, R. (2005). `spatstat`: An R package for analyzing spatial point patterns. *Journal of Statistical Software*, *12*(6). doi:[10.18637/jss.v012.i06](#)
- Hahn, T. (2005). Cuba—a library for multidimensional numerical integration. *Computer Physics Communications*, *168*(2), 78–95. doi:[10.1016/j.cpc.2005.01.010](#)
- Johnson, S. G. (2017). Adaptive multidimensional integration of a vector of integrands. Retrieved from <https://github.com/stevengj/cubature>
- Meyer, S., & Held, L. (2014). Power-law models for infectious disease spread. *Annals of Applied Statistics*, *8*(3), 1612–1639. doi:[10.1214/14-AOAS743](#)
- Meyer, S., Held, L., & Höhle, M. (2017). Spatio-temporal analysis of epidemic phenomena using the R package `surveillance`. *Journal of Statistical Software*, *77*(11). doi:[10.18637/jss.v077.i11](#)
- Narasimhan, B., Johnson, S. G., Hahn, T., Bouvier, A., & Kiêu, K. (2018). *cubature*: Adaptive multivariate integration over hypercubes. Retrieved from <https://CRAN.R-project.org/package=cubature>
- Quintero, I., Keil, P., Jetz, W., & Crawford, F. W. (2015). Historical biogeography using species geographical ranges. *Systematic Biology*, *64*(6), 1059–1073. doi:[10.1093/sysbio/syv057](#)
- R Core Team. (2018). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>
- Sommariva, A., & Vianello, M. (2007). Product Gauss cubature over polygons based on Green's integration formula. *BIT Numerical Mathematics*, *47*(2), 441–453. doi:[10.1007/s10543-007-0131-2](#)